

MAT 1275 Final Exam Review Sheet

#1 Solve for x and simplify the answer:

a) $6x^2 - 2x - 3 = 0$ b) $-2x^2 + 4x - 5 = 0$ c) $2x^2 + 2x = -3$

#2 Given the following quadratic equation, state the x and y intercepts (solutions of the form $(x, 0)$ and $(0, y)$), the vertex, and then use this information to sketch the graph:

a) $y = x^2 - 2x - 3$ b) $y = x^2 + 2x$ c) $y = -x^2 + 4x - 2$

#3 Simplify the complex fraction: a) $\frac{\frac{2}{y^2} + \frac{1}{y}}{\frac{4}{y^2} - \frac{1}{y}}$ b) $\frac{\frac{2}{x} + \frac{1}{y}}{\frac{3}{y} - \frac{4}{x}}$ c) $\frac{3 - \frac{1}{2x}}{2 + \frac{1}{x^2}}$

#4 Write the equation of circle, given below, in standard form. Identify the center and radius of the circle and graph it. Label four points on the graph with coordinates:

a) $x^2 + y^2 - 6x + 4y - 4 = 0$ b) $x^2 + y^2 + 8x - 2y - 8 = 0$
c) $x^2 + y^2 + 10x + 6y + 18 = 0$

#5 Perform the indicated operations and express the answer in the form $a + bi$:

a) $(-2 + 3i)(5 - 7i)$ b) $(4 - 9i)(3 - 2i)$ c) $\frac{3 - 2i}{1 - i}$ d) $\frac{2 - 7i}{3 + 4i}$

#6 Solve the system of equations:

$3x + 2y - z = 4$ $-2x + 5y + z = 8$ $2x - 3y + z = -9$
a) $2x - y + 2z = 10$ b) $x - 2y - 3z = -13$ c) $3x + 5y + 2z = 16$
 $x + 3y - 4z = -7$ $x + 3y - z = 5$ $-4x + 2y - 3z = 4$

#7 Solve the system of equations:

a) $3x + y = 4$
 $x^2 - 3y = -32$ b) $x^2 + y^2 = 5$
 $x - y^2 = -3$ c) $4x - y = 2$
 $x^2 + 2y = 5$ d) $x^2 - y^2 = 3$
 $2x + y^2 = 5$

8 An airplane is flying 700 feet above ground. From the plane to the base of the control tower, the angle of depression is 47° . How far away is ground directly underneath the plane to the control tower? Round your answer to the nearest tenth.

9 A temporary ramp that is 15 feet long is placed to reach an entrance door that is 4 feet above the ground. What is the angle of elevation of the ramp? Round your answer to the nearest tenth.

10 A wire, bolted to the ground 7 feet away from the base, is helping a tree stay upright. If the angle of elevation of the wire is 44° , how long is the wire? Round your answer to the nearest tenth.

11 For the given expressions

1. Identify quadrant in which angle is located
2. Find reference angle
3. Calculate the **exact** value

a) $\tan(-135^\circ)$ b) $\cos 240^\circ$ c) $\sin 330^\circ$ d) $\cos(-120^\circ)$

e) $\cot\left(-\frac{5\pi}{4}\right)$ f) $\csc\left(-\frac{4\pi}{3}\right)$ g) $\sec\left(\frac{2\pi}{3}\right)$

12 For the information given, state the value of the five remaining trig functions of θ :

a) $\sin \theta = \frac{2}{5}$ and $\cos \theta < 0$ b) $\tan \theta = 2$ and $\sin \theta < 0$

c) $\cos \theta = \frac{2}{3}$ and $\tan \theta < 0$ d) $\cos \theta = \frac{3}{7}$ and $\sin \theta < 0$

13 Given $\triangle ABC$, answer the following (round each answer to the nearest tenth):

- a) If $a = 12$, $b = 8$ and $c = 5$, find $\angle C$
- b) If $\angle A = 50^\circ$, $\angle B = 75^\circ$, and $a = 20$, find side b
- c) If $b = 9$, $c = 6$ and $\angle A = 67^\circ$, find side a
- d) If $\angle C = 37^\circ$, $\angle B = 79^\circ$, and $b = 13$, find side a

14 Prove the identities:

a) $\cos x + \sin x \cdot \tan x = \sec x$

b) $\csc x - \sin x = \cot x \cdot \cos x$

c) $\frac{1}{\cos x} - \frac{1}{\sec x} = \tan x \cdot \sin x$

d) $\csc x \cdot \cot x \cdot \sec x = 1 + \cot^2 x$

e) $\sin x \cdot \tan x \cdot \cos x = 1 - \cos^2 x$

f) $\sec x \cdot \csc x = \tan x + \cot x$

15 Find the **exact** solutions to the trig equations for $x \in [0, 2\pi)$:

a) $4 \sin x + 2 = 0$

b) $6 \cos x - 3 = 0$

c) $2 \tan x - 2 = 0$

d) $6 \tan x = -2\sqrt{3}$

16 Solve for x and round the answer to the nearest tenth:

a) $3^x = 38$

b) $2^x = 19$

c) $4^x = 7$

17 Evaluate the logarithm **without** using a calculator:

a) $\log_2 \sqrt[3]{2}$

b) $\log_3 \frac{1}{9}$

c) $\log_4 (16\sqrt[3]{4})$

d) $\log_2 (8\sqrt[6]{2})$

Answers

#1 a) $\frac{1 \pm \sqrt{19}}{6}$ b) $\frac{2 \pm i\sqrt{6}}{2}$ c) $\frac{-1 \pm i\sqrt{5}}{2}$

#2 a) Vertex: $(1, -4)$, x -intercepts: $(3, 0)$ and $(-1, 0)$, y -intercept: $(0, -3)$
 b) Vertex: $(-1, -1)$, x -intercepts: $(0, 0)$ and $(-2, 0)$, y -intercept: $(0, 0)$
 c) Vertex: $(2, 2)$, x -intercepts: $(2 - \sqrt{2}, 0)$ and $(2 + \sqrt{2}, 0)$, y -intercept: $(0, -2)$

#3 a) $\frac{2+y}{4-y}$ b) $\frac{2y+x}{3x-4y}$ c) $\frac{6x^2-x}{4x^2+2}$ or $\frac{x(6x-1)}{2(2x^2+1)}$

#4 a) Center: $(3, -2)$, Radius: $r = \sqrt{17}$ b) Center: $(-4, 1)$, Radius: $r = 5$
 c) Center: $(-5, -3)$, Radius: $r = 4$

#5 a) $11 + 29i$ b) $-6 - 35i$ c) $\frac{5}{2} + \frac{1}{2}i$ d) $-\frac{22}{25} - \frac{29}{25}i$

#6 a) $x = 3, y = -2, z = 1$ b) $x = 3, y = 2, z = 4$
 c) $x = -1, y = 3, z = 2$

#7 a) $(-5, 19), (-4, 16)$ b) $(-2, 1), (1, 2), (1, -2), (-2, -1)$
 c) $(-9, -38), (1, 2)$ d) $(2, 1), (2, -1), (-4, \sqrt{13}), (-4, -\sqrt{13})$

#8 652.8 ft **#9** 15.5° **#10** 9.7 ft

#11 a) III, 45° , 1 b) III, 60° , $-\frac{1}{2}$ c) IV, 30° , $-\frac{1}{2}$

d) III, 60° , $-\frac{1}{2}$ e) II, $\frac{\pi}{4}$, -1 f) II, $\frac{\pi}{3}$, $\frac{2\sqrt{3}}{3}$

g) II, $\frac{\pi}{3}$, -2

#12 a) $\cos \theta = -\frac{\sqrt{21}}{5}$, $\tan \theta = -\frac{2}{\sqrt{21}} = -\frac{2\sqrt{21}}{21}$, $\sec \theta = -\frac{5}{\sqrt{21}} = -\frac{5\sqrt{21}}{21}$,

$$\csc \theta = \frac{5}{2}, \cot \theta = -\frac{\sqrt{21}}{2}$$

$$\text{b) } \sin \theta = -\frac{2}{\sqrt{5}} = -\frac{2\sqrt{5}}{5}, \cos \theta = -\frac{1}{\sqrt{5}} = -\frac{\sqrt{5}}{5}, \sec \theta = -\sqrt{5},$$

$$\csc \theta = -\frac{\sqrt{5}}{2}, \cot \theta = \frac{1}{2}$$

$$\text{c) } \sin \theta = -\frac{\sqrt{5}}{3}, \tan \theta = -\frac{\sqrt{5}}{2}, \sec \theta = \frac{3}{2},$$

$$\csc \theta = -\frac{3}{\sqrt{5}} = -\frac{3\sqrt{5}}{5}, \cot \theta = -\frac{2}{\sqrt{5}} = -\frac{2\sqrt{5}}{5}$$

$$\text{d) } \sin \theta = -\frac{2\sqrt{10}}{7}, \tan \theta = -\frac{2\sqrt{10}}{3}, \sec \theta = \frac{7}{3},$$

$$\csc \theta = -\frac{7}{2\sqrt{10}} = -\frac{7\sqrt{10}}{20}, \cot \theta = -\frac{3}{2\sqrt{10}} = -\frac{3\sqrt{10}}{20}$$

#13 a) 17.6° b) 25.2 c) 8.6 d) 11.9

#14 a) $\cos x + \sin x \cdot \tan x = \cos x + \sin x \cdot \frac{\sin x}{\cos x}$
 $= \frac{\cos^2 x}{\cos x} + \frac{\sin^2 x}{\cos x} = \frac{\cos^2 x + \sin^2 x}{\cos x} = \frac{1}{\cos x} = \sec x$

b) $\csc x - \sin x = \frac{1}{\sin x} - \sin x = \frac{1}{\sin x} - \frac{\sin^2 x}{\sin x}$
 $= \frac{1 - \sin^2 x}{\sin x} = \frac{\cos^2 x}{\sin x} = \frac{\cos x}{\sin x} \cdot \cos x = \cot x \cdot \cos x$

c) $\frac{1}{\cos x} - \frac{1}{\sec x} = \frac{1}{\cos x} - \cos x = \frac{1}{\cos x} - \frac{\cos^2 x}{\cos x}$
 $= \frac{1 - \cos^2 x}{\cos x} = \frac{\sin^2 x}{\cos x} = \frac{\sin x}{\cos x} \cdot \sin x = \tan x \cdot \sin x$

d) $\csc x \cdot \cot x \cdot \sec x = \frac{1}{\sin x} \cdot \frac{\cos x}{\sin x} \cdot \frac{1}{\cos x} = \frac{1}{\sin^2 x} = \csc^2 x = 1 + \cot^2 x$

e) $\sin x \cdot \tan x \cdot \cos x = \sin x \cdot \frac{\sin x}{\cos x} \cdot \cos x = \sin^2 x = 1 - \cos^2 x$

f) For this identity, it is more convenient to start with the right hand side:

$$\begin{aligned}\tan x + \cot x &= \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{\sin^2 x + \cos^2 x}{\cos x \cdot \sin x} \\ &= \frac{1}{\cos x \cdot \sin x} = \frac{1}{\cos x} \cdot \frac{1}{\sin x} = \sec x \cdot \cos x\end{aligned}$$

#15 a) $\frac{7\pi}{6}, \frac{11\pi}{6}$ b) $\frac{\pi}{3}, \frac{5\pi}{3}$ c) $\frac{\pi}{4}, \frac{5\pi}{4}$ d) $\frac{5\pi}{6}, \frac{11\pi}{6}$

#16 a) $x = 3.3$ b) $x = 4.2$ c) $x = 1.4$

#17 a) $\frac{1}{3}$ b) -2 c) $\frac{7}{3}$ d) $\frac{19}{6}$