

# Accelerator Technologies for Simulation of Dust Impacts at Starflight Velocities

Andrew J. Higgins<sup>1\*</sup>

<sup>1</sup> McGill University, Montreal, Quebec, CANADA H3A 0C3

## Abstract

The problem of the impact of dust grains on spacecraft traveling at velocities required for interstellar travel ( $v \sim 0.1 c$ ) has been treated by only a handful of studies. [1-4] Considerable uncertainty in the physics of impacts and how the energy dissipated in the impact couples to the rest of the structure of the spacecraft exists, and it is likely that laboratory experimentation is required for validation of the computational simulation of such impacts. This paper will review existing technologies capable of accelerating macroscopic objects (i.e., impactors representative of dust grains) to ultrahigh velocities, highlighting the “no man’s land” that exists in particle mass and velocity space between launchers used in hypervelocity impacts (light gas guns, shaped-charge accelerators, electrostatic dust accelerators, electromagnetic launchers, etc.) and accelerators for fundamental particles (linacs, cyclotrons, etc.). The physics-based limitations responsible for this gap will be identified, namely (i) field-emission limits on electric field gradients for electrostatic launchers and (ii) ultrahigh current switching limitations on electromagnetic launchers. The history of the development of accelerators for impact fusion applications, likely the closest application in terms of particle mass/velocity requirements for the present problem, will be briefly reviewed. [5] Some promising—but previously unrecognized—approaches that might be able to access the particle size/velocity region of interest will be identified, such as macromolecular ion accelerators developed for mass spectrometry of biomolecules. [6] Ongoing experimental research at the author’s laboratory at McGill University to push shaped-charge accelerators capable of launching small pellets of condensed matter into the  $10^4$ - $10^6$  m/s velocity regime will also be presented. [7]

This paper is submitted for the day 2 session Sails and Beams.

**Keywords:** Accelerator Technology, Hypervelocity, Ultravelocity, Impact Physics

## References:

- [1] Benedikt, E. T. “Disintegration Barriers to Extremely High Speed Space Travel,” *Advances in the Astronautical Sciences* 6 (1961): 571-588.
- [2] Landis, G. “Dust erosion of interstellar propulsion systems,” *AIAA Paper 2000-3339, 36th AIAA/ASME/SAE/ASEE Joint Propulsion Conference and Exhibit, (2000).*
- [3] Early, J.T., and London, R.A. “Dust grain damage to interstellar laser-pushed lightsail,” *Journal of Spacecraft and Rockets* 37, no. 4 (2000): 526-531.
- [4] Hoang, T., Lazarian, A., Burkhart, B., and Loeb, A., “The Interaction of Relativistic Spacecraft with the Interstellar Medium,” *arXiv:1608.05284 (2016).*
- [5] Peaslee, A.T. Jr. (editor), *Proceedings of the Impact Fusion Workshop, LA-8000-C, Los Alamos Scientific Laboratory (1979).*
- [6] Hsu, Y.F., Lin, J.L., Lai, S.H., Chu, M.L., Wang, Y.S., and Chen, C.H., “Macromolecular ion accelerator.” *Analytical Chemistry* 84, no. 13 (2012): 5765-5769.
- [7] Georges W., Loiseau, J., Higgins, J., Tyler, T., and Zimmermann, J. “Influence of thick surface coatings on jet formation in flyer plate impacts,” *AIP Conference Proceedings* 1793, 060027 (2017).

\*Corresponding author, [andrew.higgins@mcgill.ca](mailto:andrew.higgins@mcgill.ca)